< Back

YIELD PROPHET

Save as PDF

Crop Report

12-Aug-2022

Resilient EP Soil Moisture Probe Network: Lock

Crop: Wheat

Cultivar: Scepter

Sowing details: 180 plants/m^2 on 9-May

Expected maturity date: 25-Oct

Paddock Details

Initial conditions date: 21-Mar

Soil: ResEP_sandy loam_Lock
1000 mm max rooting depth

Stubble: 1000 kg/ha of Canola
No till

Grain Yield Outcome

☑ Nitrogen limited Yield

☐ Nitrogen limited Yield with Frost and heat Effects

☑ Water limited Yield

☐ Water limited Yield with Frost and heat Effects



This graph shows the probability of exceeding a range of yield outcomes this season. It takes into account your pre-season soil moisture, the weather conditions so far, soil N and agronomic inputs. The long term record from your nominated weather station is then used to simulate what would have happened from this date on in each year of the climate record. The yield results are used to produce this graph.

Hay Yield Outcome



This graph shows the probability of exceeding a range of hay yield outcomes this season. It takes into account the same factors as the grain yield graph above. When above ground dry matter is below 2t/ha, hay yield is assumed to be 70% of dry matter, with a moisture content of 13%. When dry matter is between 2 and 12t/ha, hay yield is assumed to be between 70 and 75% of dry matter (sliding scale). When dry matter is above 12t/ha, hay yield is assumed to be between 75 and 80% (sliding scale).

Current dry matter: 6903.7kg/ha

The Season So Far - Growing Season Rainfall Deciles



Simulated and Predicted Crop Growth Stage

	GS11	GS12	GS13	GS14	GS15	GS16
	emergence	2nd leaf	3rd leaf 1st tiller	4th leaf early tillering	5th leaf mid tillering	6th leaf late tillering
Predicted						
Earliest	18-May	28-May	5-Jun	14-Jun	21-Jun	29-Jun
Median	18-May	28-May	5-Jun	14-Jun	21-Jun	29-Jun
Latest	18-May	28-May	5-Jun	14-Jun	21-Jun	29-Jun

	GS30	GS31	GS32	GS37	GS39	GS45	GS55	GS65	GS75
	end of tillering	1st node	2nd node	flag leaf	flag leaf fully emerged	mid booting	mid head emergence	mid flowering	mid dough fill
Predicted									
Earliest	17-Jul	21-Jul	26-Jul	6-Aug	11-Aug	18-Aug	27-Aug	5-Sep	22-Sep
Median	17-Jul	21-Jul	26-Jul	6-Aug	11-Aug	19-Aug	30-Aug	10-Sep	28-Sep
Latest	17-Jul	21-Jul	26-Jul	6-Aug	11-Aug	21-Aug	3-Sep	15-Sep	5-Oct

Probability and Incidence of Frost and Heat Shock

Frost damage during flowering

	Probability	This Season
mild 2 to 0°C during flowering	40%	0
moderate 0 to -2°C during flowering & early grain fill	1%	0
severe Less than -2°C during flowering & grain fill	0%	0

Heat damage during grain fill

	Probability	This Season
mild 32 to 34°C	28%	0
moderate 34 to 36°C	19%	0
severe Above 36°C	7%	0

Current Distribution of PAW



Current root depth = 1000 mm
Median final root depth = 1000 mm
Current crop PAW available to roots = 91 mm
Total Soil PAW = 103 mm
PAWC = 216 mm

PAW = Plant Available Water
CLL = Crop Lower Limit or Wilting Point
DUL = Drained Upper Limit or Field Capacity
PAWC = Plant Available Water Capacity
Current Crop PAW = Soil water currently accessible to the roots down to the current rooting depth
Soil PAW = Total accessible soil water in the soil profile

Water Budget

Initial PAW status @ 21-Mar	93 mm
Rainfall since 21-Mar	177.3 mm
Irrigations	
Evaporation since 21-Mar	86 mm
Transpiration since 21-Mar	81 mm
Deep drainage since 21-Mar	0 mm
Run-off since 21-Mar	0 mm
Current PAW status:	**103 mm**

Probability of Future Waterlogging Events



Nitrogen Budget

Initial N status @ 21-Mar	154 kg/ha
N mineralisation since 21-Mar	3 kg/ha
N tie up since 21-Mar	7 kg/ha
N applications	
	9-May : 9 kg/ha
	23-Jun : 32 kg/ha
	13-Jul : 20.7 kg/ha
Total N in plant	179 kg/ha
De-nitrification since 21-Mar	0 kg/ha
Leaching since 21-Mar	0 kg/ha
Current N status:	**32 kg/ha**

Median N mineralisation to maturity = 0.3355 kg/ha
Median N tie up to maturity = 0.373 kg/ha

Current distribution of soil nitrogen (kg/ha)



Current Crop Available N = 30 kg/ha
Total Soil N = 32 kg/ha

Availability of Water to Growing Roots



Availability of Soil Nitrogen to Growing Roots



Water Stress



1 = full stress 0 = no stress

Nitrogen Stress



1 = full stress 0 = no stress

Brief periods of mild to moderate stress do not necessarily lead to reduced yield. To see the likely impacts of additional nitrogen fertiliser rates use the Nitrogen and Nitrogen Profit reports.

Median projected crop performance and requirements for the next 10 days assuming no rain and no added fertiliser

Date	Growth Stage	Evap. (mm)	Water use (mm)	N use (kg/ha)	Water avail. to roots above stress threshold (mm)	Water avail. to roots above CLL (mm)	N avail. to roots (kg/ha)	Mineralisation (kg/ha)	N tie up (kg/ha)
13-Aug	40.8	0.4	1.6	0.6	33.4	86.8	28.0	0.0	0.0
14-Aug	41.7	0.4	1.7	0.6	31.2	84.6	27.4	0.0	0.0
15-Aug	42.6	0.4	1.5	0.5	29.4	82.7	26.9	0.0	0.0
16-Aug	43.4	0.4	1.6	0.5	27.2	80.6	26.4	0.0	0.0
17-Aug	44.2	0.4	1.6	0.5	25.0	78.4	25.9	0.0	0.0
18-Aug	45.0	0.4	1.9	0.4	23.0	76.3	25.5	0.0	0.0
19-Aug	46.0	0.4	2.0	0.4	19.8	73.2	25.1	0.0	0.0
20-Aug	46.8	0.5	2.0	0.4	17.2	70.6	24.7	0.0	0.0
21-Aug	47.8	0.5	1.9	0.4	14.7	68.1	24.3	0.0	0.0
22-Aug	48.7	0.5	1.9	0.3	12.1	65.5	24.0	0.0	0.0

The water available to roots above the stress threshold is the amount of PAW (mm) above one third of the total water holding capacity of this soil. If the water values are below this stress threshold the water available to roots above the stress threshold will be negative.

Bureau of Meteorology Seasonal and Monthly Outlooks