

Crop Report

☑Water limited Yield

13-Oct-2022 Nicole Baty: Yeelanna

Paddock Details

Initial conditions date: 24-Mar

Soil: Clay Loam over Loamy Medium Clay (Yeelanna No590) 1200 mm max rooting depth Stubble: 1000 kg/ha of Lentil No till

Grain Yield Outcome


Crop: Canola

Cultivar: Early

Sowing details: 45 plants/m² on 29-Apr


Expected maturity date: 20-Oct

☑Nitrogen limited Yield




This graph shows the probability of exceeding a range of yield outcomes this season. It takes into account your pre-season soil moisture, the weather conditions so far, soil N and agronomic inputs. The long term record from your nominated weather station is then used to simulate what would have happened from this date on in each year of the climate record. The yield results are used to produce this graph.

Hay Yield Outcome




This graph shows the probability of exceeding a range of hay yield outcomes this season. It takes into account the same factors as the grain yield graph above. When above ground dry matter is below 2t/ha, hay yield is assumed to be 70% of dry matter, with a moisture content of 13%. When dry matter is between 2 and 12t/ha, hay yield is assumed to be between 70 and 75% of dry matter (sliding scale). When dry matter is above 12t/ha, hay yield is assumed to be between 75 and 80% (sliding scale).

Current dry matter: 10451.6kg/ha



The Season So Far - Growing Season Rainfall Deciles


Simulated and Predicted Crop Growth Stage




Probability and Incidence of Frost and Heat Shock

Frost damage during flowering				Heat damage during grain fill				
Pro	bability	This Season			Pro	bability	This Season	
mild 2 to 0°C during			100%	1	mild 32 to 34°C	0%	0	
flowering			0.04	_	moderate 34 to 36°C	0%	0	
moderate 0 to -2°C during flowering & early grain fill			0%	0	Severe Above 36°C	0%	0	
SEVERE Less than -2°C during flowering & grain fill	0%	0						

Current Distribution of PAW





Current root depth = 800 mm Median final root depth = 800 mm Current crop PAW available to roots = 113 mm Total Soil PAW = 113 mm PAWC = 126 mm

PAW = Plant Available Water

- CLL = Crop Lower Limit or Wilting Point
- **DUL** Drained Upper Limit or Field Capacity
- **PAWC** = Plant Available Water Capacity
- Current Crop PAW = Soil water currently accessible to the roots down to the current rooting depth Soil PAW = Total accessible soil water in the soil profile

Water Budget		Probability of Future Waterlogging Event	ts
Initial PAW status (a) 24-Mar Rainfall since 24-Mar Irrigations Evaporation since 24-Mar Transpiration since 24-Mar Deep drainage since 24-Mar Run-off since 24-Mar Current PAW status:	90 mm 374.8 mm 166 mm 155 mm 17 mm 2 mm 113 mm	hild of the second seco	
Nitrogen Budget		Current distribution of soil nitrogen (kg/h	na)
Initial N status @ 24-Mar N mineralisation since 24-Mar N tie up since 24-Mar N applications	121 kg/ha 19 kg/ha 3 kg/ha	0 20 40 60 80	10
Total N in plant De-nitrification since 24-Mar	29-Apr : 16.1 kg/ha 16-Jun : 46 kg/ha 8-Jul : 57.5 kg/ha 243 kg/ha 1 kg/ha 4 kg/ha	500 General State 500 Control State 500 Control State Control State Control	
Leaching since 24-Mar Current N status:	4 kg/na 1 kg/ha		


Current N status:

0

May 1

Jul 1

Median N mineralisation to maturity = 0.563 kg/ha Median N tie up to maturity = 0 kg/ha



Current Crop Available N = 0 kg/ha Total Soil N = 1 kg/ha


Availability of Water to Growing Roots 500 CLL DUL Actual 400 No stress Soil water (mm)


Sep 1

Nov 1

Jan 1

Availability of Soil Nitrogen to Growing Roots





Brief periods of mild to moderate stress do not necessarily lead to reduced yield. To see the likely impacts of additional nitrogen fertiliser rates use the Nitrogen and Nitrogen Profit reports.

Median projected crop performance and requirements for the next 10 days assuming no rain and no added fertiliser

Date	Growth	Evap.	Water	N use	Water avail. to roots	Water avail. to roots	N avail.	MineralisationN tie up	
	Stage	(mm)	use (mm)	(kg/ha)	above stress threshold (mm)	above CLL (mm)	to roots (kg/ha)	(kg∕ha)	(kg/ha)
13-Oct	16.0	0.7	O.7	O.1	83.6	121.4	0.2	O.1	0.0
14-Oct	16.O	0.7	O.7	O.1	82.3	120.1	0.2	O.1	0.0
15-Oct	16.O	0.8	0.6	O.1	80.8	118.6	0.2	O.1	0.0
16-Oct	16.0	0.8	O.7	O.1	79.5	117.3	0.2	O.1	0.0
17-Oct	16.O	0.8	1.0	O.1	77.4	115.2	0.2	O.1	0.0
18-Oct	16.0	0.7	1.6	O.1	74.5	112.3	0.2	O.1	0.0
19-Oct	16.O	0.6	1.6	O.1	72.4	110.2	0.2	O.1	0.0
20-Oct	16.O	0.4	0.0	O.1	62.0	99.8	0.2	O.1	0.0
21-Oct	9.0	0.4	0.0	0.0	8.6	14.6	0.0	0.0	O.1
22-Oct	9.0	0.3	0.0	0.0	8.4	14.4	0.0	0.0	0.0

The water available to roots above the stress threshold is the amount of PAW (mm) above one third of the total water holding capacity of this soil. If the water values are below this stress threshold the water available to roots above the stress threshold will be negative.

