

Crop Report

8-Oct-2024

Andrew H Ware: Port Kenny

Crop: Wheat Cultivar: Calibre

Sowing details: 150 plants/m² on 22-Jun Expected maturity date: 2-Dec

Paddock Details

Initial conditions date: 22-Feb

Soil: Grey Calcareous Sandy Loam

(Piednippie No303)

700 mm max rooting depth Stubble: 400 kg/ha of Medic


No till

Grain Yield Outcome

♥Nitrogen limited Yield♥WaterNitrogen limited Yield with Frost and heatEffectsEffects


☑Water limited Yield

■ Water limited Yield with Frost and heat Effects



This graph shows the probability of exceeding a range of yield outcomes this season. It takes into account your pre-season soil moisture, the weather conditions so far, soil N and agronomic inputs. The long term record from your nominated weather station is then used to simulate what would have happened from this date on in each year of the climate record. The yield results are used to produce this graph.


Hay Yield Outcome




This graph shows the probability of exceeding a range of hay yield outcomes this season. It takes into account the same factors as the grain yield graph above. When above ground dry matter is below 2t/ha, hay yield is assumed to be 70% of dry matter, with a moisture content of 13%. When dry matter is between 2 and 12t/ha, hay yield is assumed to be between 70 and 75% of dry matter (sliding scale). When dry matter is above 12t/ha, hay yield is assumed to be between 75 and 80% (sliding scale).

Current dry matter: 3487.6685152139044kg/ha

The Season So Far - Growing Season Rainfall Deciles




Simulated and Predicted Crop Growth Stage




Probability and Incidence of Frost and Heat Shock

Frost damage during flowering					Heat damage during grain fill			
Probak	This Season		Probability		This Season			
mild 2 to 0°C during			0%	0	mild 32 to 34°C	54%	0	
flowering					moderate	38%	0	
moderate 0 to -2*C during flowering & early grain fill			0%	0	34 to 36°C severe Above 36°C	20%	0	
Severe Less than -2°C during flowering & grain fill	0%	0						

Current Distribution of PAW





Current root depth = 600 mm Median final root depth = 600 mm Current crop PAW available to roots = 11 mm Total Soil PAW = 11 mm PAWC = 88 mm

PAW = Plant Available Water

CLL = Crop Lower Limit or Wilting Point

DUL - Drained Upper Limit or Field Capacity

PAWC = Plant Available Water Capacity


Current Crop PAW = Soil water currently accessible to the roots down to the current rooting depth

Soil PAW = Total accessible soil water in the soil profile

Water Budget

Initial PAW status @ 22-Feb 14 mm
Rainfall since 22-Feb 187.7 mm
Irrigations
Evaporation since 22-Feb 133 mm
Transpiration since 22-Feb 127 mm
Deep drainage since 22-Feb 0 mm
Run-off since 22-Feb 0 mm
Current PAW status: 11 mm

Probability of Future Waterlogging Events




Nitrogen Budget

Initial N status @ 22-Feb 58 kg/ha N mineralisation since 22-Feb 111 kg/ha N tie up since 22-Feb 0 kg/ha N applications 21-Jun: 16 kg/ha 26-Jul: 37 kg/ha Total N in plant 90 kg/ha De-nitrification since 22-Feb 0 kg/ha Leaching since 22-Feb 0 kg/ha Current N status: 25 kg/ha


Median N mineralisation to maturity = 68.8148616018403 kg/ha Median N tie up to maturity = 0 kg/ha

Current distribution of soil nitrogen (kg/ha)




Current Crop Available N = 24 kg/ha Total Soil N = 25 kg/ha


Availability of Water to Growing Roots




Availability of Soil Nitrogen to Growing Roots



Water Stress



Nitrogen Stress



Brief periods of mild to moderate stress do not necessarily lead to reduced yield. To see the likely impacts of additional nitrogen fertiliser rates use the Nitrogen and Nitrogen Profit reports.

Median projected crop performance and requirements for the next 10 days assuming no rain and no added fertiliser

Date	Growth	Evap.	Water	N use	Water avail. to roots	Water avail. to roots	N avail.	MineralisationN tie up	
	Stage	(mm)	use	(kg/ha)	above stress threshold	above CLL (mm)	to roots	(kg/ha)	(kg/ha)
			(mm)		(mm)		(kg/ha)		
9-Oct	70.5	0.6	0.6	-0.3	-17.O	9.2	23.1	0.4	0.0
10-Oct	71.2	0.5	0.5	-0.3	-18.0	8.4	22.9	0.4	0.0
11-Oct	71.7	0.4	0.5	-0.3	-18.7	7.6	22.7	0.4	0.0
12-Oct	72.2	0.4	0.4	-0.3	-19.5	6.9	22.4	0.4	0.0
13-Oct	72.6	0.4	0.4	-0.2	-20.2	6.2	22.2	0.4	0.0
14-Oct	73.0	0.3	0.4	-0.2	-20.8	5.5	22.0	0.4	0.0
15-Oct	73.5	0.3	0.3	-0.2	-21.4	4.9	21.9	0.4	0.0
16-Oct	74.0	0.3	0.3	-0.2	-22.0	4.4	21.7	0.4	0.0
17-Oct	74.5	0.3	0.2	-0.2	-22.4	3.9	21.6	0.4	0.0
18-Oct	75.0	0.3	0.2	-0.2	-22.9	3.5	21.5	0.5	0.0

The water available to roots above the stress threshold is the amount of PAW (mm) above one third of the total water holding capacity of this soil. If the water values are below this stress threshold the water available to roots above the stress threshold will be negative.

Bureau of Meteorology Seasonal and Monthly Outlooks

