



# Crop Report

26-Jun-2024

Andrew H Ware: Port  
Kenny

Crop: Wheat

Cultivar: Calibre

Sowing details: 150 plants/m<sup>2</sup> on 22-Jun

Expected maturity date: 6-Dec

### Paddock Details

Initial conditions date: 22-Feb

Soil: Grey Calcareous Sandy Loam  
(Piednippie No303)

700 mm max rooting depth

Stubble: 400 kg/ha of Medic  
No till


## Grain Yield Outcome

Nitrogen limited Yield

Nitrogen limited Yield with Frost and heat Effects


Water limited Yield

Water limited Yield with Frost and heat Effects



This graph shows the probability of exceeding a range of yield outcomes this season. It takes into account your pre-season soil moisture, the weather conditions so far, soil N and agronomic inputs. The long term record from your nominated weather station is then used to simulate what would have happened from this date on in each year of the climate record. The yield results are used to produce this graph.


## Hay Yield Outcome



This graph shows the probability of exceeding a range of hay yield outcomes this season. It takes into account the same factors as the grain yield graph above. When above ground dry matter is below 2t/ha, hay yield is assumed to be 70% of dry matter, with a moisture content of 13%. When dry matter is between 2 and 12t/ha, hay yield is assumed to be between 70 and 75% of dry matter (sliding scale). When dry matter is above 12t/ha, hay yield is assumed to be between 75 and 80% (sliding scale).

Current dry matter: 0kg/ha

## The Season So Far - Growing Season Rainfall Deciles



# Simulated and Predicted Crop Growth Stage



## Predicted

Earliest	9-Jul	15-Jul	24-Jul	3-Aug	11-Aug	19-Aug
Median	11-Jul	19-Jul	28-Jul	6-Aug	16-Aug	24-Aug
Latest	15-Jul	22-Jul	1-Aug	12-Aug	22-Aug	1-Sep




## Predicted

Earliest	24-Aug	27-Aug	30-Aug	4-Sep	6-Sep	11-Sep	22-Sep	28-Sep	14-Oct
Median	29-Aug	2-Sep	6-Sep	14-Sep	14-Sep	22-Sep	1-Oct	6-Oct	22-Oct
Latest	3-Sep	8-Sep	14-Sep	21-Sep	22-Sep	1-Oct	10-Oct	15-Oct	2-Nov

# Probability and Incidence of Frost and Heat Shock

Frost damage during flowering				Heat damage during grain fill			
	Probability	This Season			Probability	This Season	
mild 2 to 0°C during flowering		0%	○	mild 32 to 34°C		60%	○
moderate 0 to -2°C during flowering & early grain fill		0%	○	moderate 34 to 36°C		41%	○
severe Less than -2°C during flowering & grain fill		0%	○	severe Above 36°C		26%	○

## Current Distribution of PAW



## Water Budget


Initial PAW status @ 22-Feb  
 Rainfall since 22-Feb  
 Irrigations  
 Evaporation since 22-Feb  
 Transpiration since 22-Feb  
 Deep drainage since 22-Feb  
 Run-off since 22-Feb

14 mm  
 59.1 mm  
 45 mm  
 0 mm  
 0 mm  
 0 mm

**Current PAW status:**

**29 mm**

## Probability of Future Waterlogging Events



## Nitrogen Budget

Initial N status @ 22-Feb  
 N mineralisation since 22-Feb  
 N tie up since 22-Feb  
 N applications

58 kg/ha  
 66 kg/ha  
 0 kg/ha

Total N in plant  
 De-nitrification since 22-Feb  
 Leaching since 22-Feb


21-Jun : 16 kg/ha  
 0 kg/ha  
 0 kg/ha  
 0 kg/ha

**Current N status:**

**75 kg/ha**


Median N mineralisation to maturity = 69.321995564924 kg/ha  
 Median N tie up to maturity = 0 kg/ha

## Current distribution of soil nitrogen (kg/ha)




Current Crop Available N = 24 kg/ha  
 Total Soil N = 75 kg/ha


## Availability of Water to Growing Roots




## Availability of Soil Nitrogen to Growing Roots



## Water Stress



## Nitrogen Stress




Brief periods of mild to moderate stress do not necessarily lead to reduced yield. To see the likely impacts of additional nitrogen fertiliser rates use the Nitrogen and Nitrogen Profit reports.

## Median projected crop performance and requirements for the next 10 days assuming no rain and no added fertiliser


Date	Growth Stage	Evap. (mm)	Water use (mm)	N use (kg/ha)	Water avail. to roots above stress threshold (mm)	Water avail. to roots above CLL (mm)	N avail. to roots (kg/ha)	Mineralisation (kg/ha)	N tie up (kg/ha)
27-Jun	9.0	1.1	0.0	0.0	4.6	10.1	25.9	0.3	0.0
28-Jun	9.0	0.8	0.0	0.0	4.0	9.6	26.1	0.3	0.0
29-Jun	9.0	0.6	0.0	0.0	3.7	9.2	26.3	0.4	0.0
30-Jun	9.0	0.5	0.0	0.0	3.4	9.0	26.5	0.4	0.0
1-Jul	9.0	0.5	0.0	0.0	3.2	8.8	26.8	0.4	0.0
2-Jul	9.0	0.4	0.0	0.0	3.1	8.7	27.0	0.4	0.0
3-Jul	10.0	0.4	0.0	-0.1	3.0	8.7	27.6	0.4	0.0
4-Jul	10.1	0.4	0.0	-0.1	3.1	9.5	28.4	0.4	0.0
5-Jul	10.3	0.3	0.0	0.0	3.1	10.3	30.9	0.4	0.0
6-Jul	10.4	0.3	0.0	0.0	3.2	11.2	33.6	0.4	0.0

The water available to roots above the stress threshold is the amount of PAW (mm) above one third of the total water holding capacity of this soil. If the water values are below this stress threshold the water available to roots above the stress threshold will be negative.


### 3 MONTH RAINFALL OUTLOOK FOR OCTOBER TO DECEMBER




OCTOBER RAINFALL OUTLOOK




NOVEMBER RAINFALL OUTLOOK



### PAST ACCURACY FOR OCTOBER TO DECEMBER



PAST ACCURACY FOR OCTOBER



PAST ACCURACY FOR NOVEMBER

